

N.S. Bala, M.D., F.A.C.G., F.A.C.G., A.G.A.F. S. K. Raj, M.D., F.A.C.P.,F.A.S.G.E. H.S Ojeas, M.D., F.E. Scheinder, M.D., D.H. Darmadi, M.D., F.A.C.G., F.A.S.G.E. J. Rao, M.D., F.A.C.G., M.I. Kafrouni, M.D. A.H. Ghazale., M.D., J.P.Shah, M.D. Diplomates, American Board of Internal Medicine & Gastroenterology

OsmoPrep® INSTRUCTIONS

DO NOT TAKE ASPIRIN, ANTI-INFLAMMATORY MEDICATION, BLOOD THINNERS, OR IRON SUPPLEMENTS ONE WEEK PRIOR TO YOUR PROCEDURE

DAY BEFORE PROCEDURE:

- 1. Stay on a Clear Liquid Diet all day (see back for details →) AVOID ANYTHING WITH RED COLOR DYE. It is extremely important that you drink plenty of fluids to prevent dehydration.
- 2. At 6 p.m. begin taking 4 OsmoPrep tablets every 15 minutes with at least 8 ounces of any clear liquid (for a total of 20 tablets).
- 3. NOTHING TO EAT OR DRINK AFTER MIDNIGHT.

DAY OF PROCEDURE:

- 1. 5 hours prior to your procedure begin taking 4 OsmoPrep tablets every 15 minutes with at least 8 ounces of any clear liquid (for a total of 12 tablets).
- * You may take important medicines such as heart medication or blood pressure medication with a small sip of water.
- * FOR DIABETICS ONLY: Reduce your Insulin dose by _ the day prior to your procedure. DO NOT take Insulin the morning of the procedure until after you have eaten. Please clarify instructions with the physician who has prescribed the Insulin.
- * You will be sedated. You **MUST** have someone drive you home. You should also have someone with you for 4 to 6 hours after you are home. Family members or driver will need to stay at the facility until you have been discharged. Instructions will be given to them after your procedure.
- * Please wear comfortable clothes and shoes the day of your procedure.

PROCEDURE:	
PLACE:	DR:
DATE:	<u> </u>
PROCEDURE TIME:	<u> </u>
Check in atAM/PM	
☐ At Outpatient Registration on the	floor.
☐ At Bay Area Endoscopy Center	

NOTE: There will be a hospital or facility charge for any procedure in addition to the physician charge.